



POST-OPERATIVE INSTRUCTIONS FOR DENTAL IMPLANT PATIENTS

These instructions will guide you through the first few days of recovery.

All patients heal differently - listen to your body. If something hurts, do not do it!

Start taking the prescribed antibiotics and rinse (2x per day for 1 minute) 3 days before surgery.

Bleeding is Normal and Expected

- After leaving the office, gently bite on gauze, replacing it with new gauze as needed. Remove gauze when eating or drinking, then replace if bleeding persists.
- If bleeding is not controlled with gauze alone, moisten a tea bag then apply it directly to the extraction site. This will help your body create a blood clot.
- Blood (pink) tinged saliva is normal for the first 24-48 hours.

Numbness Will Last for Several Hours After Surgery

- Begin taking post-operative medications shortly after you arrive home. Follow the medication instructions on the reverse page.
- After numbness has subsided you may add other warm nourishing foods to your diet sticking with soft foods that require limited chewing. This includes soups, scrambled eggs, and mashed potatoes.
- Drinking through a straw IS permitted.

Dietary Guidelines

- Eat soft, cold nourishing food limiting your intake to liquids, pureed or soft-cooked foods. High calorie and high protein meal replacement shakes are recommended to have on hand.

Swelling is Normal and Expected

- Swelling will typically peak about 24-48 hours after surgery and will subside within a week.
- During the first 24-48 hours, we recommend applying ice packs to help control bleeding.

- Once the bleeding is under control and swelling begins, you may switch to warm, moist heat to help with discomfort. (Gel packs can be warmed in the microwave for 10 seconds)

Sutures Will Dissolve Over the Next 7-10 Days

- But may come out as early as the ride home if you move your mouth too much. They cannot be replaced.

Restrict Your Physical Activity for 5–7 Days After Surgery

- You may return to work or school as you feel ready, typically 2-3 days after surgery.

Oral Hygiene is Important and Promotes Healing

- If you smoke or Vape, do not smoke during the first 48 hours.
- Rinse gently with warm salt water 3-4 times per day for the next 7-10 days, beginning the morning after surgery.
- Brush your teeth and avoid surgical areas.

IMPLANT INSTRUCTIONS

- An implant healing abutment is a temporary cap that may have been placed on the implant. The healing abutment will be visible in the mouth and is not the implant but a “cover” that is threaded into the implant. A healing abutment is used to guide the gum tissue and allow it to heal in the shape of the implant.
- If the healing abutment comes loose or falls off during the healing period, please call our office and we will replace or tighten the healing abutment. A loose or lost healing abutment will not damage the implant if it is addressed properly.

The local anesthetic administered with the general anesthetic during your surgery will last for 3-6 hours after surgery.

What Can I Do For Pain?

Some discomfort is normal after surgery.

To minimize pain and swelling take **600mg (3 tablets) Ibuprofen** (ex. Advil or Motrin) every 8 hours for the first 3-4 days post surgery combined with **Tylenol Extra Strength (500mg)** Take the first dose before the anesthesia wears off.

If prescription pain medication is given, take it as instructed on the label. Do not exceed the dose on the label.

Taking medications with food or milk will help reduce upset stomach.

Avoid driving or operating heavy machinery when taking prescription pain medication.

Do not drink alcohol while taking prescription pain medications.

MEDICATIONS: Along with pain medication you will be prescribed antibiotics as indicated by the procedure. Take the antibiotics as prescribed until the entire prescription is completed

IF YOU ARE HAVING ANY PROBLEMS, PLEASE CALL US 313-884-3050 AS SOON AS POSSIBLE.